

レジメン名	(胃癌) イリノテカン+シスプラチン High dose	適応														胃癌													
		1クール日数														28日													
		上限クール数														なし													
投与量\day		1	2	3	4	5	6	7	8					14	15							21	22					28	
【0.0】mL/瓶	1 瓶	●													●														
液【3.3】mg・1mL/管 液【6.6】mg・2mL/瓶 75mg・5mL/瓶 【0.0】mL/瓶	1 管 1 瓶 1 瓶 1 瓶														●														
【0.0】mL/瓶	1 瓶	●													●														
塩点滴静注液「タイリ」	70mg/m2														●														
500mL/袋	1 袋	●													●														
射（薬剤調製）																													
00mL/袋	1 袋	●																											
ン点滴静注「マルコ」	80mg/m2																												
】mL/袋(点滴用)	1 袋	●																											
射（薬剤調製）																													
注射液10mg・2mL/管 00mL/袋	2 管 1 袋	●																											
00mL	1 袋	●																											
00mL	1 袋	●																											
注 500mL (ガイーンF)	1 瓶		●																										
注 500mL (ガイーンD) て	1 瓶		●																										
00mL	1 袋		●																										
【0.0】mL/瓶	1 瓶														●														
		1	2	3	4	5	6	7	8					14	15								21	22					28
mg タ 食 後	2 錠		●	●	●																								
カプセル 125mg の60～90分前	1 C	●																											
カプセル 80mg 食 後	1 C		●	●																									

胃/化療/10h≤

胃/化療/≤3h