

レジメン名	(大腸癌) パニツムマブ+イリノテカン 2週間隔投与	適応	大腸癌
		1クール日数	35日
		上限クール数	なし

【注射】		投与量\day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	
<<主管①>> 生理食塩液【100】mL/瓶 キープ用 点滴注射	1 瓶		●														●																					
<側管①> ■ベクティビックス点滴静注用(100) 生理食塩液【100】mL/瓶 60分かけて ベクティビックス投与後1時間経過観察 抗癌剤点滴注射(薬剤調製)	6mg/kg 1 袋		●														●																					
<側管②> 生理食塩液【50】mL/瓶 経過観察用 30分かけて 点滴注射	1 瓶		●														●																					
<側管③> デキサート注射液【3.3】mg・1mL/管 デキサート注射液【1.65】mg・0.5mL/管 アロキシ静注0.75mg・5mL/瓶 生理食塩液100mL/瓶 30分かけて 点滴注射	1 管 1 管 1 瓶 1 瓶		●														●																					
<側管④> 生理食塩液【50】mL/瓶 フラッシュ用 点滴注射	1 瓶		●														●																					
<側管⑤> ■イリノテカン塩酸塩点滴静注液「タイウ」 (40/100) テルモ生食【500】mL/袋(点滴用) 90分かけて 抗癌剤点滴注射(薬剤調製)	150mg/m ² 1 袋		●														●																					
<側管⑥> 生理食塩液【50】mL/瓶 フラッシュ用 点滴注射	1 瓶		●														●																					
フィルター付輸液セット			●														●																					
【内服】			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	
アプレピタントカプセル 125mg 1日1回 点滴の60~90分前	1 C		●														●																					
アプレピタントカプセル 80mg 1日1回 朝 食後	1 C			●	●													●	●																			

腸/化療/≦4h