レジメン名		適応	大腸癌			
レンバンロ	(大腸癌)ベバシズマブ+SOX	1クール日数	2 1 日			
		上限クール数	なし			

														上队	見ク	<u> </u>	ル数	Į		な	し	
F. S. 413	10 L E	_			-	_		_														
【注射】 《主管① ≫	投与量\day	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
【≪王官①グ 【生理食塩液【100】mL/瓶	1 瓶																					
キープ用	. 7120																					
点滴注射																						
<側管①>	7.5 //																					
■アバスチン点滴静注用 (100 /400)	7.5mg/kg																					
(100/400) 生理食塩液【100】mL/瓶	1 瓶																					
90分かけて	1 TEL	•																				
※初回の忍容性が良ければ2回目6																						
0分3回目以降30分で投与可能。																						
抗癌剤点滴注射(薬剤調製)																						
<側管②>									Г													
生理食塩液【50】m L/瓶	1 瓶																					
フラッシュ用																						
点滴注射 <側管①>																						
〜 側 音 ① <i>/</i> デキサート注射液【6.6】mg⋅2mL/瓶	1 瓶																					
ラモセトロン注射液 0.3mg·2mL/管	1 管																					
生理食塩液【100】mL/瓶	1 瓶	-																				
30分かけて																						
点滴注射																						
<側管②> ■オキサリプラチン注(50/100	130mg/m²																					
デキサート注射液【3.3】mg·1mL/管	1 管																					
大塚糖液【5% 250mL】/瓶	1 瓶	•																				
2時間かけて																						
抗癌剤点滴注射(薬剤調製)																						
<側管③> 生理食塩液【50】mL/瓶	l 1 瓶																					
プラッシュ用	1 TEL																					
点滴注射																						
																						_
【内服】		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
■エスワンタイホウ配合0D錠T 1日2回 朝 夕 食 後	™ g	タ [^]	Ţ						П						1	~ ~朝						
デカドロン錠 4mg	2 錠	ŕ														+7.						_
1日2回朝 夕 食後	- "-		•	•																		
																					, ,	
																			胄/'	比療	!/≦	4h
	(エスワン																					
		体表面積 初回基準量																				
	└─ ┤					~	-1 .	25r							ıg/			〈 2				
			1. 2	5m2			~ 1			未清	5				ıg/			< 2				
	(1. 3	om2	じ	上	·~				б	Οn	ıg/	口	(>	〈 2)			